



## Metal impact in the marine system

Dong, Yan; Hauschild, Michael Zwicky

*Publication date:*  
2011

[Link back to DTU Orbit](#)

*Citation (APA):*

Dong, Y., & Hauschild, M. Z. (2011). *Metal impact in the marine system*. Poster session presented at 3rd NorLCA Symposium, Helsinki, Finland.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.




# Metal impact in the marine system

Yan Dong & Michael. Z. Hauschild


Section for Quantitative sustainability Assessment (QSA), Department of Management Engineering  
Technical University of Denmark (DTU), Kgs, Lyngby, Denmark

E-mail: yado@man.dtu.dk

## Introduction



Small River



Metals are used during the life cycle of mobile phones. Most of the metals are recycled in EU, but emissions from these facilities may reach soil and water. In cases where effective recycling infrastructures are missing, e.g. in developing countries, this also occurs when the end of life treatment is inadequate.

The emissions to water will finally end up in the marine system, resulting in potential risks to marine organisms.

### Aim of the study:

Among all metal species, only the free metal ions are toxic to marine organisms. Taking Cu and Zn as examples, this study presents how metals are transported in the marine water, and differentiate into different species. The metals' toxic risks are revealed.

## Materials and methods

### Toxic metal speciation:

$$\text{Free metal ion } (M^{2+}) = (M_T) * (M^{2+}/M_T)$$

$M_T$  is the total dissolved concentration

A small continuous discharge source (discharge rate  $10\text{m}^3/\text{s}$ ) emitting metal-polluted wastewater into a middle sized bay (Donegal Bay) is simulated by MIKE21<sup>#</sup>. In addition to hydrodynamic modeling, total dissolved metal ( $M_T$ ) concentrations are simulated simultaneously.

Fraction  $M^{2+}/M_T$  is estimated based on empirical data obtained from various literature sources.

<sup>#</sup>MIKE21 is a model of coast and sea, developed by Danish Hydraulic Institute

## Results




Fig. 1.  $\text{Cu}_T$  concentration (mg/l) after 2 months discharge

Due to precipitation and adsorption to the sediments and particles, both Cu and Zn concentrations decrease to less than 10% of their source concentrations after 5km. When reaching the edge of the bay, they further decrease to less than 1%.




Fig. 2.  $\text{Zn}_T$  concentration (mg/l) after 2 months discharge




Fig. 3. Empirical  $\text{Cu}^{2+}/\text{Cu}_T$

Fig. 3 shows that in the marine water, concentration of  $\text{Cu}^{2+}$  is at least 2 orders of magnitudes lower than  $\text{Cu}_T$ , which means that  $\text{Cu}^{2+}/\text{Cu}_T < 1\%$ . Fig. 4 shows that  $\text{Zn}^{2+}$  is 0.5—2 magnitudes lower than  $\text{Zn}_T$ , which means that  $1\% < \text{Zn}^{2+}/\text{Zn}_T < 32\%$ .

Combined with Fig. 1. and Fig. 2., when reaching bay's edge,  $\text{Cu}^{2+}/(\text{Cu}_T \text{ at source}) < 0.01\%$ , which results in  $1.5 \mu\text{g/l Cu}^{2+}$ ;  $\text{Zn}^{2+}/(\text{Zn}_T \text{ at source}) < 0.32\%$ , which results in  $16 \mu\text{g/l Zn}^{2+}$ . Comparing to free ion EC50s of marine organisms, which are  $120.6 \mu\text{g/l}$  for  $\text{Cu}^{2+}$  and  $263.2 \mu\text{g/l}$  for  $\text{Zn}^{2+}$  (data calculated from epa.org database), the emissions probably will not cause adverse effects to the marine organisms.




Fig. 4. Empirical  $\text{Zn}^{2+}/\text{Zn}_T$

## Conclusion

From this study, it appears that most of the metal emissions precipitate to the sediments before they enter the marine system, or they are bound to ligands in the marine water reducing their availability to marine life. Only a small portion of metals in the marine system exist as toxic free ions, which may cause very limited damages to the marine organisms. Therefore, metals impacts in the marine environment is highly overestimated by existing ecotoxicity characterization methods for use in LCA which consider the full dissolved fraction as toxic and tend to disregard the strong attenuation in estuaries.